

ORIGINAL ARTICLE

Inhibition of blood cholinesterases by nerve agents *in vitro*

Jiří Bajgar¹, Daniel Jun^{1,2}, Kamil Kuča^{1,2}, Josef Fusek¹, Jana Žďárová Karasová¹, Jiří Kassa¹, Jiří Cabal¹, Václav Bláha³

¹Department of Toxicology, ²Center of Advanced Studies, ³Department of Multidisciplinary Studies, Faculty of Military Health Sciences, University of Defence, Hradec Králové, Czech Republic

Received 17th June 2009.

Revised 8th September 2009.

Published online 12th November 2009.

Summary

The bimolecular rate constants (k_a) of inhibition were determined for cholinesterase in rat blood *in vivo* and three nerve agents *in vitro*. They were in the range of 10^5 – 10^6 M·min⁻¹ with the highest value for VX ($k_a = 8.92 \times 10^6$), the next highest for soman ($k_a = 3.22 \times 10^6$) and the lowest for sarin ($k_a = 0.39 \times 10^6$), respectively. The inhibition rate is the same *in vitro* and *in vivo*; when the real inhibition *in vivo* and *in vitro* is compared, it is possible to assess the concentration of the nerve agent present in the blood stream. When administered intramuscularly it is about 70% of the dose administered.

Key words: whole blood cholinesterases; rat; nerve agents; inhibition *in vitro*

INTRODUCTION

The toxicodynamics of nerve agents and organophosphates (OP) are explained by acetylcholinesterase (AChE, EC 3.1.1.7) inhibition at the cholinergic synapses and the consequent accumulation of neuromediator acetylcholine followed by metabolic disbalance and other changes (Marrs 1996, Eyer 2003, Bajgar 2004, Rotenberg and Newmark 2004, Wiener and Hoffmann 2004, Geoghegan 2006). Exposure to OP is characterized by four basic reactions in the organism: resorption, transport, metabolism and toxic effect (Bajgar 2004).

Before the toxic action, the OP is transported by the blood stream to the target sites (the central and peripheral nervous systems). The transport system can be considered as a base for the further action of the OP, and it is the first main system on which the OP have an effect. This reaction is very important because a part of the OP is bound and this part of the administered dose is thus excluded from further toxic action at the target sites, as is a further part that in some compounds is detoxified: thus, G agents (sarin, soman etc.) are, and V agents (VX, Russian VX etc.) are not detoxified in the organism (Bajgar 1991).

In the transport system (blood), two cholinesterases are present; AChE in the erythrocytes and butyrylcholinesterase (BuChE, EC 3.1.1.8) in the plasma. Both enzymes are inhibited during the transport of the OP according to the principle of “first come, first served” (Benschop and de Jong 2001), and therefore this inhibition can be considered as a good marker of poisoning with cholinesterase inhibitors (Bajgar et al. 2006). The partitioning between these two enzymes is influenced by the affinity of AChE and BuChE to the nerve agents. There are other

✉ Jiří Bajgar, Department of Toxicology,
Faculty of Military Health Sciences,
University of Defence, Třebešská 1575,
500 01 Hradec Králové, Czech Republic

💻 bajgar@pmfhk.cz

☎ +420 973 251 507

📠 +420 495 518 094

enzymes and proteins binding the nerve agents: carboxylesterases (inhibited by the agents) and blood proteins – particularly albumin – (binding the agents), with, however, significantly lower affinity than nerve agents (Bajgar 1991, 2004). The inhibition of both cholinesterases in blood depends mostly on:

- the affinity of OP to the cholinesterases in blood;
- the rate of penetration of the OP from the route of administration to the transport system;
- the detoxification of the OP.

The affinity of OP to cholinesterases can easily be tested *in vitro*, and this approach is frequently evaluated in the research of new reactivators. Although there are many studies on the inhibition and reactivation of cholinesterases *in vitro* (Kuča et al. 2004, 2005, Patočka et al. 2005, Musilek et al. 2007, Petroianu and Kalasz 2007, Hasan et al. 2008, Petroianu and Lorke 2008), there is a lack of complex data on the most usual and available animal – the rat – for all tissues of interest. There are separate data on the affinity *in vitro* of different OP to: human (Jandorf et al. 1955, Tammelin 1957, Aquilonius et al. 1964, Patočka and Tulach 1969, Aurbek et al. 2006); pig (Bajgar et al. 1971, Aurbek et al. 2006); erythrocyte AChE, and plasma BuChE (Tammelin 1957, Aquilonius et al. 1964, Back 1969, Bajgar et al. 1971), or brain AChE (Back 1969, Patočka and Bajgar 1971, Bajgar and Patočka 1977) etc. Inhibition constants inform us about the kinetics of the enzymes mentioned but do not correspond to the real status, i.e. to the whole inhibition capacity in the blood. However, this information is necessary for comparison of *in vivo* and *in vitro* results. The aim of this study is to obtain inhibition constants characterizing the affinity of the most common nerve agents – sarin, soman and VX – to the blood cholinesterases (plasma BuChE and erythrocyte AChE determined as the activity in the whole blood), *in vitro* using one method (one substrate, pH, conditions etc.) and *in vivo* (the rat).

MATERIAL AND METHODS

Animals

6 female Wistar rats of the same litter (BioTest, Konárovice, Czech Republic), weighing 200–220 g, were used. The animals were housed in the Central Vivarium of the Faculty of Military Health Sciences under veterinary control. All the experiments were performed with the permission and under the supervision of the Ethics Committee of the Faculty of Military Health Sciences, Hradec Králové (permission No 153/06) according to § 17 of Czech law No. 207/2004, permission of responsible person

0001/94 – M 699. The animals were exsanguinated and the heparinized blood was used for the experiments.

Chemicals

Nerve agents were obtained from the Military Technical Institute of Protection (Brno, Czech Republic). They were of minimally 98% purity and stored in glass ampullas (1 ml). The solutions for experiments were prepared immediately before use.

Determination of cholinesterase activity/inhibition

Inhibition of the blood samples was performed using a solution of the relevant nerve agent (sarin, soman, VX) in isopropanol, a mixture of the whole blood (495 µl) and the inhibitor solution (5 µl). The final concentration of the inhibitor was 10^{-6} , 10^{-7} a 10^{-8} M, and the final concentration of the solvent was 1% (v/v). At different time intervals (0, 3, 6, 10, 15, 20, 30 min), an aliquot part of the blood sample (50 µl) was haemolysed by adding of 950 µl distilled water. After 1 min, AChE activity was determined according to Ellman et al. (1961) as follows: 100 µl of the haemolysate was mixed with 1700 µl DTNB solution (1 mM solution of DTNB in 0.1 M TRIS-HCl buffer, pH 7.6) and an enzymatic reaction was started by adding 200 µl of substrate solution (1 mM acetylthiocholine in distilled water). The mixture was stabilized for 3 min and then the absorbancy was monitored at 436 nm/min (the delay is necessary to exclude false positive results caused by the titration of free SH-groups interfering with the activity determination).

Statistical evaluation

The dependence of cholinesterase activity changes vs. time was evaluated by the least square method in semi-logarithmical transformation, using relevant PC programmes: relative activity (Δ absorbancy_{436 nm}) was recalculated to per cent values (activity at the time zero represents 100%). Evaluation of the data was carried out using Microsoft Excel 2002 and GraphPad Prism version 4 for Windows, GraphPad Software, San Diego, California, USA (www.graphpad.com). For each determination (time interval, concentration of inhibitor), the mean from 3 measurements was used.

RESULTS

Absolute values of cholinesterase activity in the rat blood varied from 10.1 to 15.5 µkat/l, with an average 11.3 ± 1.12 µkat/l (1 kat = $60 \cdot 10^6$ unit; unit/l = 16.67 nkat/l).

Changes of the activity in the blood after incubation with sarin are shown in Fig. 1. The inhibition is dependent on the inhibitor concentration and the time of incubation. Semilogarithmic transformation of the data allows a determination of the half-life of inhibition as shown in Fig. 2.

The bimolecular rate constants (k_a) were calculated according to the equation

$$\frac{0.69}{[I] \cdot t_{0.5}} = k_a \text{ (equation I);}$$

the same approach was used for calculation of k_a values for soman and VX. They are summarized in Table 1. According to the bimolecular rate constants, VX seems to be the most potent inhibitor of rat blood cholinesterases followed by sarin and soman. From these data, it is possible to calculate the IC_{50} constant, i.e. the inhibitor concentration causing 50% of inhibition. The values for 10 min inhibition are also shown in Table 1. It is clear that the order of IC_{50} values is the same (sarin < soman < VX) but they are given for easier comparison with the data in the literature.




Fig. 1. Inhibition of cholinesterases in the rat blood by different concentrations of sarin *in vitro*. Sarin concentrations: 0.01–0.1 and 1.0 μM .




Fig. 2. Inhibition of blood cholinesterases by different concentrations of sarin *in vitro* in semilogarithmic transformation. Sarin concentrations: 0.01–0.1 and 1.0 μM .




Fig. 3. Correlation between toxicity expressed as LD_{50} (i.m., $\mu\text{g/kg}$) and bimolecular rate constants for sarin, soman, VX (our data), and O-ethyl 2-(2-dimethylaminoethyl) methyl phosphonothiolate (data of Bajgar 1991) in semilogarithmic and linear transformation. The correlation coefficients (R^2) are different for different transformations; higher R^2 for semilogarithmic transformation indicates that this dependence will be probably more significant.

DISCUSSION

The inhibition of erythrocyte AChE of different species has been described many times, and the erythrocyte source has mostly been human, pig or rat; e.g. Aurbek et al. (2006) determined the bimolecular rate constant (k_a) for human erythrocyte AChE to be $9.91 \cdot 10^7$ and for pig erythrocyte AChE and VX, $4.43 \cdot 10^7 \text{ M} \cdot \text{min}^{-1}$. These constants are higher than those obtained in our experiments: this could be caused by the use of a different enzyme source and different contents of AChE and BuChE in the blood. This ratio for rat blood (7:3) was determined in an earlier study (71% for erythrocyte AChE and 29% for plasma BuChE) (Bajgar 1972). The sensitivity of blood cholinesterases to sarin and VX has been described – for human erythrocyte AChE – as about 10 times higher for VX in comparison with sarin (Worek et al. 2004). Jandorf et al. (1955) determined the bimolecular rate constant for rat erythrocyte AChE to be approximately 10^7 , and this is in agreement with our data. The inhibition constant occurring in the literature more frequently – IC_{50} – can be derived from the equation (I), but this constant is very dependent on the time of incubation of the enzyme with an inhibitor. Among other tissues studied for AChE inhibition, the data for human

Table 1. Inhibition constants for interaction of the rat blood cholinesterases with nerve agents.

Constant	Sarin	Soman	VX
$k_a \cdot 10^6 \text{ (M} \cdot \text{min}^{-1}\text{)}$	0.39 ± 0.26	3.22 ± 0.88	8.92 ± 1.03
$I_{50} \cdot 10^{-7} \text{ (M)}$	176	22.4	7.7
LD ($\mu\text{g/kg}$, i.m.)*	200 (180)	70 (77)	13.6 (15)

* Toxicity data according to Bajgar (2004) and the reference Vachek et al. (1996).

erythrocyte AChE activity are as follows: Back (1969) determined IC_{50} for human plasma and soman 10^{-7} – 10^{-8} M, whereas the same constant for pig erythrocyte AChE was 10^{-9} M (Bajgar et al. 1971) and for human plasma BuChE 10^{-8} M (Tammelin 1957, Aquilonius et al. 1964, Patočka and Tulach 1969, Bajgar and Patočka 1977). The correlation between toxicity and bimolecular rate constants for sarin, soman, VX (our data), and O-ethyl 2-(2-dimethylaminoethyl) methyl phosphonothiolate (Bajgar 1992) is demonstrated in Fig. 3.

If we compare the inhibition potency characterized by k_a constant with the toxicity expressed as LD_{50} (Vachek et al. 1996, Bajgar et al. 2007), a good correlation can be found (Fig. 3). In all these comparisons it is necessary to point out that inhibition of cholinesterases in the blood is an indicator of exposure but it is not connected with the mechanism of the action of nerve agents: it has been demonstrated that inhibition of erythrocyte AChE in rabbits does not influence normal functions of the organism (Kassa et al. 1997, Kassa and Bajgar 1999). However, the inhibition in the transport system can be very good marker of not only exposure level but also a marker for assessment of nerve agent concentration in the blood stream. This was our purpose in making further calculations:

It was demonstrated that following administration of VX (i.m., $1 \cdot LD_{50}$, $13.6 \mu\text{g/kg}$, i.e. $5.087 \cdot 10^{-8}$ mol/kg), the half life ($t_{0.5}$) of blood cholinesterase inhibition was determined to be 2.4 min (Bajgar et al. 2007). When this constant is introduced into the equation (I), the concentration of inhibitor (VX) was calculated as $3.667 \cdot 10^{-8}$ mol/l/kg, and it is 72.1% of the dose administered. This means that after i.m. administration, about 70% of the dose administered is present in the blood stream. Using this approach, it would be possible to calculate a dose of inhibitor present in the blood after intoxication with different routes of administration, and thus, to improve our knowledge of the mechanism of the action of these highly toxic agents.

CONCLUSIONS

For studies dealing with the effect of nerve agents, whole blood is the material of choice for inhibition studies.

For kinetic studies, the use of separated enzymes is recommended.

The inhibition of cholinesterases in whole blood allows an assessment of the concentration of inhibitor in the blood stream.

Toxicity correlates well with the rate of cholinesterase inhibition in the blood.

ACKNOWLEDGEMENT

The authors are indebted to Mrs J. Uhlířová, M. Hrabínová and M. Zechovská for skillful technical assistance. Financial support of the Ministry of Defence, grant No FVZ 0000501 is gratefully acknowledged.

REFERENCES

- Aquilonius SM, Fredriksson T, Sundwall A: Studies on phosphorylated thiocholine and choline derivatives. I. General toxicology and pharmacology. *Toxicol Appl Pharmacol* 6:269–279, 1964.
- Aurbek N, Thiermann H, Szinicz L, Eyer P, Worek F: Application of kinetic-based computer modelling to evaluate the efficacy of HI-6 in percutaneous VX poisoning. *Toxicology* 224:74–80, 2006.
- Back P: Zur Reaktivierung der Azetylcholinesterase und Cholinesterase. *Titrimetrische Untersuchungen an Hirnhomogenat und Plasma nach Tabun-, Sarin- und Somanhemmung in vitro*. *Z Klin Chem Klin Biochem* 7:301–305, 1969.

- Bajgar J: Time course of acetylcholinesterase inhibition in the medulla oblongata of the rat by O-ethyl S-(2-dimethylaminoethyl) methylphosphonothioate *in vivo*. *Brit J Pharmacol* 45:368-371, 1972.
- Bajgar J: The influence of inhibitors and other factors on cholinesterases. *Sb Ved Pr Lek Fak Karlovy Univerzity (Hradec Králové)* 34:3-75, 1991.
- Bajgar J: Biological monitoring of exposure to nerve agents. *Brit J Ind Med* 49:648-653, 1992.
- Bajgar J: Organophosphates/nerve agent poisoning: mechanism of action, diagnosis, prophylaxis and treatment. *Adv Clin Chem* 38:151-216, 2004.
- Bajgar J, Patočka J: *In vitro* inhibition of soluble brain acetylcholinesterase by organophosphates of O-ethyl-S-(2-dialkylaminoethyl)-methylphosphonothiolate type. *Collect Czech Chem Commun* 42:770-772, 1977.
- Bajgar J, Vondráček V, Sajda M: Inhibition of blood cholinesterases of the pig by some organophosphates *in vitro* and *in vivo* (in Czech). *Vojen Zdrav Listy* 40:266-270, 1971.
- Bajgar J, Fusek J, Bartošová L, Jun D, Kuča K: Evaluation of reactivation test in anaesthetized dogs with experimental intoxication with nerve agents. *J Appl Toxicol* 26:439-443, 2006.
- Bajgar J, Kuča K, Fusek J, Karasová J, Kassa J, Cabal J, Jun D, Bláha V: Inhibition of blood cholinesterases following intoxication with VX and its derivatives. *J Appl Toxicol* 27:458-463, 2007.
- Benschop HP, de Jong LPA: Toxicokinetics of nerve agents. In Somani SM, Romano RM (eds): *Chemical Warfare Agents: Toxicity at Low Levels*. CRC Press, Boca Raton 2001, pp 25-81.
- Ellman GL, Courtney DK, Andres V, Featherstone RM: A new and rapid colorimetric determination of acetylcholinesterase activity. *Biochem Pharmacol* 7:88-95, 1961.
- Eyer P: The role of oximes in the management of organophosphorus pesticides poisoning. *Toxicol Rev* 22:165-190, 2003.
- Geoghegan T: Chemical warfare agents. *Anaesth Crit Care Pain* 6:230-234, 2006.
- Hasan MY, Lorke DE, Arafat K, Kuča K, Musilek K, Schmitt A, Petroianu GA: *In vitro* oxime protection of red blood cell acetylcholinesterase inhibited by diisopropyl-fluorophosphate. *Clin Toxicol* 46:397-397, 2008.
- Jandorf JB, Michel HO, Schaffer NK, Egan R, Sommerson WH: The mechanism of reaction between esterases and phosphorus containing antiesterases. *Discuss Faraday Soc* 20:134-142, 1955.
- Kassa J, Bajgar J, Fusek J: Changes of cholinesterase activity in the erythrocytes, plasma, diaphragm, liver and various parts of the brain in the rabbit following transfusion of erythrocytes with soman inhibited acetylcholinesterase. *Acta Med (Hradec Králové)* 40:37-39, 1997.
- Kassa J, Bajgar J: *In vitro* inhibition of rabbit and guinea pig erythrocyte acetylcholinesterase by soman. *Toxicol In Vitro* 13:403-408, 1999.
- Kuča K, Pícha J, Cabal J, Liška F: Synthesis of the three monopyridinium oximes and evaluation of their potency to reactivate acetylcholinesterase inhibited by nerve agents. *J Appl Biomed* 2:51-56, 2004.
- Kuča K, Cabal J, Bajgar J, Jun D: *In-vitro* screening for a new potent reactivator of acetylcholinesterase inhibited by a nerve agent VX. *Lett Drug Des Discov* 2:23-25, 2005.
- Marrs TC: Organophosphate anticholinesterase poisoning. *Toxic Subst Mech* 15:357-388, 1996.
- Musilek K, Kuča K, Jun D, Doležal M: *In vitro* reactivation potency of bispyridinium (E)-but-2-ene linked acetylcholinesterase reactivators against tabun-inhibited acetylcholinesterase. *J Appl Biomed* 5:25-30, 2007.
- Patočka J, Bajgar J: Affinity of human brain acetylcholinesterase to some organophosphates and carbamates *in vitro*. *J Neurochem* 18:2545-2546, 1971.
- Patočka J, Tulach J: Interaction of O-ethyl S-(2-dimethylaminoethyl) methylthiophosphonate with acetylcholinesterase. *Collect Czech Chem Commun* 34:3191-3195, 1969.
- Patočka J, Cabal J, Kuča K, Jun D: Oxime reactivation of acetylcholinesterase inhibited by toxic phosphorus esters: *in vitro* kinetics and thermodynamics. *J Appl Biomed* 3:91-99, 2005.
- Petroianu GA, Kalasz H: Comparison of the ability of pyridinium aldioximes to reactivate human RBC cholinesterases inhibited by ethyl- and methyl-paraoxon. *Curr Med Chem* 11:1624-1634, 2007.
- Petroianu GA, Lorke DE: Pyridinium oxime reactivators of cholinesterase inhibited by diisopropyl-fluorophosphate (DFP). Predictive value of *in vitro* testing for *in vivo* efficacy. *Mini Rev Med Chem* 8:1328-1342, 2008.
- Rotenberg JS, Newmark J: Nerve attacks on children: diagnosis and management. *Pediatrics* 112:648-658, 2004.
- Tammelin LP: Dialkoxy-phosphorylthiocholines, alkoxy-methylphosphorylthiocholines and

- analogous choline esters. Synthesis, pKa of tertiary homologues and cholinesterase inhibition. *Acta Chem Scand* 11:1340–1348, 1957.
- Vachek J, Geršl V, Fusek J, Krs O, Skopec F, Bajgar J: Toxicities of O-alkyl S-(2-dialkylaminoethyl) methyl phosphonothiolates. *Acta Med (Hradec Králové)* 39:67–71, 1996.
- Wiener SW, Hoffman RS: Nerve agents: a comprehensive review. *J Int Care Med* 19:22–37, 2004.
- Worek F, Thiermann, H, Szinicz L, Eyer P: Kinetic analysis of interactions between human acetylcholinesterase, structurally different organophosphorus compounds and oximes. *Biochem Pharmacol* 68:2237–2248, 2004.